# Clinical guideline PICU early mobilization

Pediatric intensive care unit

# PICU early mobilization guideline criteria

## Level 1

- Intubated, FIO2 >60%, PEEP >8\*
- Difficult/critical airway, new trach/prior to first trach change
- Acute neurological event, elevated ICP
- Sedated and RASS score > -3 or < +2
- Active titration of vasoactive drugs
- Unstable pelvis/spine without stabilization device
- **CONTRAINDICATIONS:** active or uncontrolled bleeding/seizures, or open chest/abdomen

# Level 2

- Intubated/or trach <60% and/or PEEP <8\*
- Non-invasive support with FIO2 >60%\*
- Dialysis/CRRT
- Femoral Access

## Level 3

- Non-invasive pulmonary support with FIO2 <60%</li>
- Baseline pulmonary support
- EVD clamped by nurse and RASS -1 to +3

\*consider baseline ventilator settings

# Children's Hospital of Richmond at VCU

# PICU early mobilization guideline activity

Level 1

- Lights on, shades up by 0900; dimmed by 2300
- Consider low stimulation (noise machines, limited TV, etc.)
- HOB to/ > 30 degrees
- Turn q2h during day and consider q4 at night
- OT consult by PICU day 3
- PT as needed
- Developmentally supportive positioning per PT and OT
- PROM initiated as appropriate

#### Level 2

## **Rest and reassess**

- Change in HR 20%
- Change in BP 20%
- Change in RR 20%
- Decrease SpO2 15%
- Increase FiO2 20%
- Increase ETCO2 20%
- Ventilator asynchrony
- CPAP/BiPAP asynchrony
- Respiratory distress
- New arrhythmia
- Hemodynamic concerns
- Change in mental status
- Concern for airway device, vascular access, or EVD integrity
- Behavior interfering with safe activity

- Level 1 activities PLUS:
- Positive touch for infants/toddlers
- Sitting up in bed TID
- Team to consider OOB to chair and/or ambulation
- PT and OT consult by day 3
- Assess for difficulty with communication, phonation, and swallowing readiness for SLP consult
- Assess need for daily schedule
- CAPD BID for delirium screen

Level 3

- Level 1 and 2 activities PLUS:
- OOB to chair TID or sitting up in bed if appropriate chair not available
- Ambulation BID with assist

For questions concerning this guideline, contact: chorclinicalguidelines@vcuhealth.org chrichmond.org Last updated: January 2023 Next expected update: January 2026

# PICU early mobilization guideline **Executive summary**

## Pediatric intensive care unit early mobilization guideline workgroup

Pediatric physical therapy owner: Jennifer Resetar, PT DPT Pediatric occupational therapy owner: Jacqueline Reissis, OT Pediatric critical care owner: Colleen Wisnewski, RN Pediatric critical care: Alia O'Meara, MD Pediatric critical care: Tammie Shatzer, RN QI expert: Michelle Olson, MD Pediatric occupational therapy owner: Tamara Neiderer, COTA Pediatric critical care owner: Dale Purrington, CPNP-AC Pediatric critical care: Nikki Miller Ferguson, MD Pediatric critical care: Kara Greenfield, MD Respiratory therapy: Michelle Humrich, RT

# Approved (May 2023)

Medical director of pediatric critical care medicine Mark Marinello, MD

**CHoR clinical guidelines committee:** Jonathan Silverman, MD, MPH; Ashlie Tseng, MD **CHoR quality council, executive sponsor:** Elizabeth Peterson RN, MPH, CPHQ

Matthew Schefft, DO, MSHA

# **References:**

#### Guideline team members

- 1. Lee CM, Fan E. ICU-acquired weakness: what is preventing its rehabilitation in critically ill patients? BMC Med. 2012 Oct 3;10:115.
- Moraes FDS, Marengo LL, Moura MDG, Bergamaschi CC, de Sá Del Fiol F, Lopes LC, Silva MT, Barberato-Filho S. ABCDE and ABCDEF care bundles: A systematic review of the implementation process in intensive care units. Medicine (Baltimore). 2022 Jun 24;101(25):e29499.
- 3. Smith HAB, Besunder JB, Betters KA, Johnson PN, Srinivasan V, Stormorken A, Farrington E, Golianu B, Godshall AJ, Acinelli L, Almgren C, Bailey CH, Boyd JM, Cisco MJ, Damian M, deAlmeida ML, Fehr J, Fenton KE, Gilliland F, Grant MJC, Howell J, Ruggles CA, Simone S, Su F, Sullivan JE, Tegtmeyer K, Traube C, Williams S, Berkenbosch JW. 2022 Society of Critical Care Medicine Clinical Practice Guidelines on Prevention and Management of Pain, Agitation, Neuromuscular Blockade, and Delirium in Critically III Pediatric Patients With Consideration of the ICU Environment and Early Mobility. Pediatr Crit Care Med. 2022 Feb 1;23(2):e74-e110.
- Wieczorek B, Ascenzi J, Kim Y, Lenker H, Potter C, Shata NJ, Mitchell L, Haut C, Berkowitz I, Pidcock F, Hoch J, Malamed C, Kravitz T, Kudchadkar SR. PICU Up!: Impact of a Quality Improvement Intervention to Promote Early Mobilization in Critically III Children. Pediatr Crit Care Med. 2016 Dec;17(12):e559-e566.

